



Marioni, R. E., Suderman, M., Chen, B. H., Horvath, S., Bandinelli, S., Morris, T., Beck, S., Ferrucci, L., Pedersen, N. L., Relton, C. L., Deary, I. J., & Hägg, S. (2018). Tracking the Epigenetic Clock Across the Human Life Course: A Meta-analysis of Longitudinal Cohort Data. *Journals of Gerontology, Series A*.
<https://doi.org/10.1093/gerona/gly060>

Peer reviewed version

Link to published version (if available):
[10.1093/gerona/gly060](https://doi.org/10.1093/gerona/gly060)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Oxford University Press at <https://academic.oup.com/biomedgerontology/advance-article/doi/10.1093/gerona/gly060/4944478> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Tracking the epigenetic clock across the human life-course - a meta-analysis of longitudinal cohort data

Riccardo E. Marioni, PhD^{1,2}, Matthew Suderman, PhD³, Brian H. Chen, PhD⁴, Steve Horvath, PhD^{5,6}, Stefania Bandinelli, MD⁷, Tiffany Morris, PhD⁸, Stephan Beck, PhD⁸, Luigi Ferrucci, MD, PhD⁴, Nancy L. Pedersen, PhD⁹, Caroline L. Relton, PhD³, Ian J. Deary, PhD^{1,10}, and Sara Hägg, PhD^{9,*}

1. Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, 7 George Square, Edinburgh, EH8 9JZ, UK
2. Medical Genetics Section, Centre for Genomic and Experimental Medicine, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, EH4 2XU, UK
3. Medical Research Council Integrative Epidemiology Unit, University of Bristol, Bristol, UK
4. The National Institute on Aging, NIH, Baltimore, MD, USA
5. Human Genetics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA 90095, USA
6. Department of Biostatistics, School of Public Health, University of California Los Angeles, Los Angeles, CA 90095, USA
7. Geriatric Unit, Azienda Sanitaria di Firenze, Florence, Italy
8. UCL Cancer Institute, University College London, London, UK
9. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, 17177 Stockholm, Sweden
10. Department of Psychology, University of Edinburgh, 7 George Square, Edinburgh, EH8 9JZ, UK

*Corresponding author:

Sara Hägg, PhD

Department of Medical Epidemiology and Biostatistics

Karolinska Institutet

Box 281

171 77 Stockholm, Sweden

Sara.hagg@ki.se

Target journal: The Journals of Gerontology: Series A

Word count: Brief report (2800 words excluding title page and abstract), max 20 refs

Abstract (250 words)

Background

Epigenetic clocks based on DNA methylation yield high correlations with chronological age in cross-sectional data. Due to a paucity of longitudinal data, it is not known how Δ_{age} (epigenetic age – chronological age) changes over time or if it remains constant from childhood to old age. Here, we investigate this using longitudinal DNA methylation data from five datasets, covering most of the human life course.

Methods

Two measures of the epigenetic clock (Hannum and Horvath) are used to calculate Δ_{age} in the following cohorts: ALSPAC offspring (n~950, total age-range 7-19 years, 2 waves), ALSPAC mothers (n~900, 16-60 years, 2 waves), InCHIANTI (n~460, 21-100 years, 2 waves), SATSA (n~370, 48-99 years, 5 waves), Lothian Birth Cohort 1936 (n~1,050, 70-76 years, 3 waves), and Lothian Birth Cohort 1921 (n~470, 79-90 years, 3 waves). Linear mixed models were used to track longitudinal change in Δ_{age} within each cohort.

Results

For both epigenetic age measures, Δ_{age} showed a declining trend in almost all of the cohorts. The correlation between Δ_{age} across waves ranged from 0.22 to 0.82 for Horvath and 0.25 to 0.71 for Hannum, with stronger associations in samples collected closer in time.

Conclusions

Epigenetic age increases at a slower rate than chronological age across the life course, especially in the oldest population. Some of the effect is likely driven by survival bias, where healthy individuals are those maintained within a longitudinal study, although other factors

like the age distribution of the underlying training population may also have influenced this trend.

Key words: epigenetic clock, longitudinal, life-course perspective

Introduction

A number of studies have demonstrated age-related methylation differences at specific CpG sites. Indeed, linear combinations of CpG methylation beta values – labelled epigenetic clocks – correlate highly with chronological age (Pearson $r > 0.90$) (1, 2). For a given chronological age, older epigenetic age is presumed to indicate poorer health, and has been associated with increased mortality risk (3) and many age-related morbidities (4, 5).

Because published epigenetic clocks were derived from cross-sectional data, it is unknown whether individual differences between epigenetic age and chronological age (Δ_{age}) are (i) set at birth and continue unchanged over the life course, (ii) changing gradually across the life course or (iii) changing more notably during specific periods of life e.g., adolescence and old age. Such questions can be tested using cross-sectional data, although repeated measurements at multiple times hold clear advantages for inference, especially because they are not biased by selective survival. In this study we use longitudinal methylation data from five population-based cohorts, spanning the life course from early childhood to death.

Methods

Longitudinal DNA methylation data were collected in five cohorts: the Avon Longitudinal Study of Parents and Children (ALSPAC), Invecchiare in Chianti (InCHIANTI), the Swedish Adoption/Twin Study of Aging (SATSA), the Lothian Birth Cohort 1936 (LBC1936), and the Lothian Birth Cohort 1921 (LBC1921).

ALSPAC is a “transgenerational prospective observational study investigating influences on health and development across the life course”(6, 7). Participants comprise a cohort of offspring born to pregnant women recruited in 1991-2 in Bristol, UK. Participants have been followed through a series of ongoing data collection waves involving questionnaires and clinical assessments. Please note that the study website contains details of all the data that is available through a fully searchable data dictionary

(<http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary/>). DNA methylation was measured in the peripheral blood of the offspring at ages 7 and 15-17 (986 individuals corresponding to 1,901 samples) and of their mothers during pregnancy and approximately 18 years later (982 individuals corresponding to 1,816 samples). The resulting profiles form part of the Accessible Resource for Integrated Epigenomics Studies (ARIES) dataset (8). Data are available by request from the Avon Longitudinal Study of Parents and Children Executive Committee (<http://www.bristol.ac.uk/alspac/researchers/access/>).

The **InCHIANTI** study is a population-based prospective cohort study of residents aged 20 or older from two areas in the Chianti region of Tuscany, Italy. Sampling and data collection procedures have been described elsewhere (9). Briefly, 1,326 participants donated a blood sample at baseline (1998-2000), of which 784 also donated a blood sample at the 9-year follow-up (2007-2009). DNA methylation was assayed in participants with sufficient DNA at both baseline and year 9 visits (n=499). After samples and data quality checks, DNA methylation data at baseline and follow-up were available in 460 individuals.

The **SATSA** study is a longitudinal prospective cohort study of adult Swedish twins (10, 11). It was started in 1984 and has been ongoing until 2014. There are up to ten waves of in-person testing available with questionnaire data on health and life-style choices, cognitive

testing, physical performance measures, anthropometrics, and blood draws. DNA methylation was assessed repeatedly up to five times in 373 individuals corresponding to 938 samples. The age ranges in the methylation SATSA samples spanned from 48 to 89 years at baseline (1992) and 63 to 99 years at the last follow-up (2012).

LBC1921 and **LBC1936** are birth cohorts containing Scottish participants born in 1921 and 1936, most of whom participated in the Scottish Mental Surveys of 1932 and 1947, when nearly all 11-year old Scottish children completed a cognitive test (12, 13). Longitudinal follow up of those living in the Lothian area began in 2000 for those born in 1921 and in 2006 for those born in 1936. Blood-based DNA methylation data were available at three times in both cohorts – mean ages 70, 73, and 76 years in LBC1936 (1,054 individuals corresponding to 2,338 samples), and mean ages 79, 87, and 90 in LBC1921 (476 individuals corresponding to 703 samples).

Ethics

In ALSPAC, informed written consent was obtained from parents of participants after receiving a complete description of the study at the time of enrolment into the ALSPAC project, with the option for them or their children to withdraw at any time. Ethical approval for the ALSPAC study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees.

InCHIANTI participants provided written informed consent to participate in this study. The study complied with the Declaration of Helsinki. The Italian National Institute of Research

and Care on Aging Institutional Review Board approved the study protocol and study participants provided informed consent.

Participants in SATSA provided informed consents at each testing occasion. The longitudinal collection and analyses of data have been approved at several occasions by the Research Ethics Committee at Karolinska Institutet with Dnrs 84:61, and 98-319, and by the Regional Ethics Board in Stockholm with Dnrs 2007/151-31/4, 2010/657-31/3, and 2015/1729-31/5.

Following written informed consent, venesected whole blood was collected for DNA extraction in both LBC1921 and LBC1936. Ethics permission for the LBC1921 was obtained from the Lothian Research Ethics Committee (Wave 1: LREC/1998/4/183). Ethics permission for the LBC1936 was obtained from the Multi-Centre Research Ethics Committee for Scotland (Wave 1: MREC/01/0/56), the Lothian Research Ethics Committee (Wave 1: LREC/2003/2/29), and the Scotland A Research Ethics Committee (Waves 2 and 3: 07/MRE00/58).

DNA Methylation and Epigenetic Clock Measurements

Blood-based Illumina 450k methylation data were obtained separately in each cohort. Cohort specific quality control details have been reported previously (14-17). Epigenetic age was calculated by multiplying beta values by the regression weights from Horvath (2) and Hannum et al (1) to create the respective clocks. The Hannum clock was derived using DNA methylation from blood in a single cohort of 656 individuals; the Horvath clock was derived using DNA methylation from 51 tissue types across 8,000 individuals from multiple studies. The Hannum clock was developed from the Illumina 450K array, while the Horvath clock was restricted to approximately 21,000 probes common to both the Illumina 27K and 450K

arrays. Further, the Hannum clock is moderately correlated with proportions of certain blood cells, while the Horvath clock is relatively uncorrelated with blood cell counts to date. Delta age was defined as a simple subtraction of chronological age from epigenetic age using both versions of the clock. Cell count predictions were estimated from DNA methylation data using the Houseman method in all cohorts (18).

Statistical Analysis

Linear mixed models were used to assess longitudinal change in Δ_{age} separately in each cohort. Δ_{age} was modelled as the outcome, with chronological age as the time-scale and predictor of interest. All models controlled for sex and a random effect intercept term. The SATSA study further adjusted for the twin structure in the data by allowing for additional random effects within twin pairs. Analyses were conducted in R using the lme4 and lmerTest packages. Pearson correlations were calculated for Horvath and Hannum Δ_{age} between waves for all pair-wise combinations in each cohort. Sensitivity analyses were carried out using only individuals present in at least three waves and using cell count prediction adjustments for the Hannum clock.

Results

The total number of individuals participating in this study was 4,075 with a total of 8,616 samples. Summary statistics of chronologic and epigenetic age at each study wave of the participating cohorts are presented in Table 1. Individuals from the different cohorts represent most of the human life course, from young children to the oldest old, although the majority of the samples were drawn in later life. All cohorts with the exception of the ALSPAC mothers

had an even distribution of men and women. The number of follow-up occasions in the cohorts varied from two to five, with different time intervals in between the measurements.

The fitted mean trajectories of Δ_{age} over time for each cohort are presented together in Figure 1. As the Hannum clock was not trained on children, the ALSPAC children were excluded from the Hannum plot (Figure 1B). Participant level trajectories for each cohort for the Horvath- and Hannum epigenetic clock measures, as well as respective Δ_{age} , are shown in Supplementary Figure 1A-D. Judging from the trajectories and the mixed model output (Supplementary Table 1), Δ_{age} declines during adulthood (β_{Horvath} ranged from -0.18 to -0.40, all $P < 2 \times 10^{-16}$; β_{Hannum} ranged from -0.09 to -0.62, all $P < 4 \times 10^{-16}$), and also during childhood, although to a smaller degree ($\beta_{\text{Horvath}} = -0.07$, $P = 8 \times 10^{-7}$). In other words, over time, the DNA methylation-based biological clock increases at a slower rate than chronological age. The only exception to this trend was the Lothian Birth Cohort 1936, where Δ_{age} remained constant over a 6-year interval between ages 70 and 76 years ($\beta_{\text{Horvath}} = -0.01$, $P = 0.81$; $\beta_{\text{Hannum}} = -0.04$, $P = 0.16$). Sensitivity analyses keeping only individuals with at least three waves of measurements ($n = 183$ in SATSA, 487 in LBC1936, and 66 in LBC1921) did not change the results (Supplementary Table 2). Likewise, adjustments for predicted cell counts for the Hannum epigenetic clock measurements did not change the age estimates remarkably (Supplementary Table 3).

The correlation between Δ_{age} across waves is presented by cohort in Table 2, and ranged from 0.22 to 0.82 for Horvath and 0.25 to 0.71 for Hannum. The association of between-wave correlations and sampling times between the waves is illustrated in Figure 2, where increasing sampling time confers a lower correlation between samples for Horvath Δ_{age}

(Beta=-0.015 units per year, p-value= 9.3×10^{-4}), but less so for Hannum Δ_{age} (Beta=-0.009 units per year, p-value=0.039). Sensitivity analyses keeping only individuals with three measures confirmed the decreasing correlation pattern for Horvath Δ_{age} (Beta=-0.016 units per year, p-value= 2.5×10^{-4}) but not for Hannum Δ_{age} (Beta=-0.005 units per year, p-value=0.28).

Discussion

In this paper we presented the first comprehensive analysis of the epigenetic clock from a longitudinal perspective by analysing data from five prospective cohorts with repeated sampling. We showed that epigenetic age was highly correlated with chronologic age when including multiple samples per individual, and that Δ_{age} declined over the life span.

Moreover, cross-correlations of Δ_{age} from different waves indicated more similar patterns in samples collected closer in time compared to samples collected further apart, although the pattern was more prominent in Horvath than in Hannum estimates.

The epigenetic clock has been shown to be a useful marker of biological age when using data from cross-sectional sample collections (19). Here, we provided evidence for its usefulness in a longitudinal perspective. The overall correlations with chronologic age were high, as judged from the individual trajectory plots, and thus comparable to cross-sectional study findings. However, the trajectories of Δ_{age} showed a declining trend in almost all of the cohorts with adult sample collections. This indicates that epigenetic age increases at a slower rate than chronological age, especially in the oldest population. Some of the effect is likely driven by survival bias, where healthy individuals are those maintained within a longitudinal study, although other factors like underlying training population for the respective clocks may also

have influenced this trend. It may also be possible that there is a ceiling effect for Δ_{age} whereby epigenetic clock estimates plateau. In children, the Horvath epigenetic age declined with chronologic age, although less so than in the adult life span, while the Hannum clock was not trained in children and hence was not used.

The investigation of correlations between different waves in each cohort showed a decreasing correlation for samples collected further apart, which is in line with expectations where measures further apart have been more influenced by other (environmental) factors.

However, it should be noted that the trend is somewhat different for Horvath and Hannum clocks. This is perhaps not surprising given that they represent different tissues; Horvath is a multi-tissue clock built to capture more variation while Hannum only applies to blood leukocytes (only a few CpG sites overlap in the two clocks) and all our samples came from blood leukocytes.

The strength of this study is the joint effort of combining five longitudinal cohorts, comprising six data sets, with repeated sample collections assessed by DNA methylation 450k arrays. By doing so, we were able to capture the full life-course perspective of the epigenetic clock from childhood to old age. However, there is an overrepresentation of samples collected at the later part of the life span, which limits the interpretations. Moreover, these cohorts are all based on individuals from a European ancestry background. As there is evidence for differences based on the epigenetic clock in other ethnic populations (20), our findings are not necessarily generalizable to other ethnicities.

In summary, we have provided an analysis of longitudinal trajectories of the epigenetic clock across the life-course, showing that epigenetic age increases at a slower rate than chronological age across the life course, especially in the oldest population

Funding

The UK Medical Research Council and Wellcome (grant number 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. This publication is the work of the authors and Matthew Suderman will serve as guarantor for the ALSPAC-related contents of this paper. Analysis of the ALSPAC data was funded by UK Economic & Social Research Council grant (grant number ES/N000498/1). ARIES was funded by the BBSRC (BBI025751/1 and BB/I025263/ 1). Supplementary funding to generate DNA methylation data which are (or will be) included in ARIES has been obtained from the MRC, ESRC, NIH and other sources. ARIES is maintained under the auspices of the MRC Integrative Epidemiology Unit at the University of Bristol (grant numbers MC_UU_12013/2, MC_UU_12013/8 and MC_UU_12013/9).

The InCHIANTI study baseline (1998-2000) was supported as a "targeted project" (ICS110.1/RF97.71) by the Italian Ministry of Health and in part by the U.S. National Institute on Aging (Contracts: 263 MD 9164 and 263 MD 821336).

The SATSA study was supported by NIH grants R01 AG04563, AG10175, AG028555, the MacArthur Foundation Research Network on Successful Aging, the Swedish Council for Working Life and Social Research (FAS/FORTE) (97:0147:1B, 2009-0795, 2013-2292), the Swedish Research Council (825-2007-7460, 825-2009-6141, 521-2013-8689, 2015-03255),

KI Foundation, the Strategic Research Program in Epidemiology at Karolinska Institutet, and by Erik Rönnbergs donation for scientific studies in aging and age-related diseases.

REM and IJD conducted the research in The University of Edinburgh Centre for Cognitive Ageing and Cognitive Epidemiology (CCACE), part of the cross-council Lifelong Health and Wellbeing Initiative (MR/K026992/1); funding from the Biotechnology and Biological Sciences Research Council (BBSRC) and Medical Research Council (MRC) is gratefully acknowledged. Phenotype collection in the Lothian Birth Cohort 1921 was supported by the UK's Biotechnology and Biological Sciences Research Council (BBSRC), The Royal Society and The Chief Scientist Office of the Scottish Government. Phenotype collection in the Lothian Birth Cohort 1936 was supported by Age UK (The Disconnected Mind project). Methylation typing was supported by Centre for Cognitive Ageing and Cognitive Epidemiology (Pilot Fund award), Age UK, The Wellcome Trust Institutional Strategic Support Fund, The University of Edinburgh, and The University of Queensland.

Acknowledgements

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses.

Figure legends

Figure 1. Mean linear longitudinal trajectories of epigenetic Δ_{age} . For each data set, mixed models were applied and predicted values, derived from the model intercept and fixed effect estimates for age, were plotted to illustrate the Δ_{age} trajectories across the life span. The x-axis represents the age where the cohort specific trajectory is plotted corresponding to the age span covered in that cohort. The y-axis shows the Δ_{age} .

Figure 2. Within-cohort correlations of epigenetic Δ_{age} across study waves. For each data set, correlations between all possible combinations of waves were calculated and plotted. On the x-axis is the time between the two measurements in years and on the y-axis is the correlation coefficient.

References

1. Hannum G, Guinney J, Zhao L, Zhang L, Hughes G, Sada S, *et al.* Genome-wide methylation profiles reveal quantitative views of human aging rates. *Mol Cell.* 2013;**49**:359-367; doi:10.1016/j.molcel.2012.10.016.
2. Horvath S. DNA methylation age of human tissues and cell types. *Genome Biol.* 2013;**14**:R115; gb-2013-14-10-r115 [pii]; doi:10.1186/gb-2013-14-10-r115.
3. Chen BH, Marioni RE, Colicino E, Peters MJ, Ward-Caviness CK, Tsai PC, *et al.* DNA methylation-based measures of biological age: meta-analysis predicting time to death. *Aging (Albany NY).* 2016;**8**:1844-1865; doi:10.18632/aging.101020.
4. Horvath S, Ritz BR. Increased epigenetic age and granulocyte counts in the blood of Parkinson's disease patients. *Aging (Albany NY).* 2015;**7**:1130-1142;
5. Marioni RE, Shah S, McRae AF, Ritchie SJ, Muniz-Terrera G, Harris SE, *et al.* The epigenetic clock is correlated with physical and cognitive fitness in the Lothian Birth Cohort 1936. *Int J Epidemiol.* 2015; dyu277 [pii]; doi:10.1093/ije/dyu277.
6. Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, *et al.* Cohort Profile: the 'children of the 90s'--the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol.* 2013;**42**:111-127; doi:10.1093/ije/dys064.
7. Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Davey Smith G, *et al.* Cohort Profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol.* 2013;**42**:97-110; doi:10.1093/ije/dys066.
8. Relton CL, Gaunt T, McArdle W, Ho K, Duggirala A, Shihab H, *et al.* Data Resource Profile: Accessible Resource for Integrated Epigenomic Studies (ARIES). *Int J Epidemiol.* 2015;**44**:1181-1190; doi:10.1093/ije/dyv072.
9. Ferrucci L, Bandinelli S, Benvenuti E, Di Iorio A, Macchi C, Harris TB, *et al.* Subsystems contributing to the decline in ability to walk: bridging the gap between epidemiology and geriatric practice in the InCHIANTI study. *J Am Geriatr Soc.* 2000;**48**:1618-1625;
10. Finkel D, Reynolds CA, McArdle JJ, Gatz M, Pedersen NL. Latent growth curve analyses of accelerating decline in cognitive abilities in late adulthood. *Dev Psychol.* 2003;**39**:535-550;
11. Pedersen NL, McClearn GE, Plomin R, Nesselroade JR, Berg S, DeFaire U. The Swedish Adoption Twin Study of Aging: an update. *Acta Genet Med Gemellol (Roma).* 1991;**40**:7-20;
12. Deary IJ, Gow AJ, Pattie A, Starr JM. Cohort profile: the Lothian Birth Cohorts of 1921 and 1936. *Int J Epidemiol.* 2012;**41**:1576-1584; doi:10.1093/ije/dyr197.
13. Deary IJ, Gow AJ, Taylor MD, Corley J, Brett C, Wilson V, *et al.* The Lothian Birth Cohort 1936: a study to examine influences on cognitive ageing from age 11 to age 70 and beyond. *BMC Geriatr.* 2007;**7**:28; doi:10.1186/1471-2318-7-28.
14. Shah S, McRae AF, Marioni RE, Harris SE, Gibson J, Henders AK, *et al.* Genetic and environmental exposures constrain epigenetic drift over the human life course. *Genome Res.* 2014;**24**:1725-1733; doi:10.1101/gr.176933.114.
15. Wang Y, Karlsson R, Lampa E, Zhang Q, Hedman ÅK, Almgren M, *et al.* Epigenetic influences on aging: a longitudinal genome-wide methylation study in old Swedish twins. *bioRxiv.* 2017; doi:10.1101/226266.
16. Dedeurwaerder S, Defrance M, Calonne E, Denis H, Sotiriou C, Fuks F. Evaluation of the Infinium Methylation 450K technology. *Epigenomics.* 2011;**3**:771-784; doi:10.2217/epi.11.105.
17. Moore AZ, Hernandez DG, Tanaka T, Pilling LC, Nalls MA, Bandinelli S, *et al.* Change in Epigenome-Wide DNA Methylation Over 9 Years and Subsequent Mortality: Results From the InCHIANTI Study. *J Gerontol A Biol Sci Med Sci.* 2016;**71**:1029-1035; doi:10.1093/gerona/glv118.
18. Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, *et al.* DNA methylation arrays as surrogate measures of cell mixture distribution. *BMC Bioinformatics.* 2012;**13**:86; doi:10.1186/1471-2105-13-86.

19. Jylhava J, Pedersen NL, Hagg S. Biological Age Predictors. *EBioMedicine*. 2017; 10.1016/j.ebiom.2017.03.046.
20. Horvath S, Gurven M, Levine ME, Trumble BC, Kaplan H, Allayee H, *et al*. An epigenetic clock analysis of race/ethnicity, sex, and coronary heart disease. *Genome Biol*. 2016;**17**:171; doi:10.1186/s13059-016-1030-0.

Table 1. Characteristics of longitudinal methylation cohorts

Cohort	Wave	Year	Participants (N)	Women (%)	Age mean \pm SD	Horvath Age mean \pm SD	Hannum Age mean \pm SD
SATSA	1	1992-1994	212	60%	68.3 \pm 9.1	60.3 \pm 9.9	65.3 \pm 9.3
	2	1999-2001	227	63%	71.0 \pm 10.1	63.2 \pm 8.8	66.9 \pm 8.9
	3	2002-2004	178	54%	72.1 \pm 9.2	64.0 \pm 8.8	68.3 \pm 9.1
	4	2008-2010	172	61%	75.9 \pm 8.2	67.3 \pm 9.2	71.0 \pm 7.8
	5	2010-2012	149	66%	77.8 \pm 8.2	67.3 \pm 9.1	71.4 \pm 7.8
LBC1936	1	2004-2007	920	49%	69.6 \pm 0.8	66.0 \pm 6.5	71.3 \pm 5.8
	2	2007-2010	800	48%	72.5 \pm 0.7	69.3 \pm 6.6	72.9 \pm 5.7
	3	2011-2013	618	48%	76.3 \pm 0.7	72.6 \pm 6.4	77.6 \pm 5.6
LBC1921	1	1999-2001	446	60%	79.1 \pm 0.6	73.7 \pm 7.0	80.3 \pm 6.2
	2	2007-2008	175	54%	86.7 \pm 0.4	77.6 \pm 6.0	81.6 \pm 6.2
	3	2011-2012	82	54%	90.1 \pm 0.9	79.3 \pm 6.1	84.8 \pm 5.8
ALSPAC Children	1	1998-2000	948	50%	7.5 \pm 0.15	8.3 \pm 2.4	9.2 \pm 4.6
	2	2006-2010	953	52%	17.1 \pm 1.0	17.2 \pm 4.3	20.4 \pm 4.9
ALSPAC Mothers	1	1991-1992	924	100%	29.1 \pm 4.4	30.2 \pm 6.7	34.9 \pm 5.6
	2	2008-2011	892	100%	47.4 \pm 4.5	45.1 \pm 6.5	47.1 \pm 6.2
InCHIANTI	1	1998-2000	460	54%	62.2 \pm 16.1	61.6 \pm 13.3	67.6 \pm 16.0
	2	2007-2009	460	54%	71.4 \pm 16.2	68.7 \pm 13.1	75.2 \pm 15.9

Table 2. Pearson correlations of epigenetic Δ_{age} between waves in each cohort

SATSA

Wave	1	2	3	4	5
1		0.51	0.51	0.61	0.46
2	0.48		0.61	0.56	0.62
3	0.44	0.46		0.67	0.58
4	0.33	0.51	0.53		0.63
5	0.22	0.40	0.51	0.43	

LBC1936

Wave	1	2	3
1		0.49	0.49
2	0.61		0.65
3	0.56	0.69	

LBC1921

Wave	1	2	3
1		0.54	0.57
2	0.64		0.71
3	0.55	0.64	

ALSPAC children

Wave	1	2
1		0.31
2	0.26	

ALSPAC mothers

Wave	1	2
1		0.25
2	0.28	

InCHIANTI

Wave	1	2
1		0.71
2	0.82	

White: Horvath; Grey: Hannum